Zero Excess and Minimal Length in Finite Coxeter Groups

S.B. Hart and P.J. Rowley*

Abstract

Let $\mathcal{W}$ be the set of strongly real elements of W, a Coxeter group. Then for $w \in \mathcal{W}$, $e(w)$, the excess of w, is defined by $e(w) = \min\{\ell(x) + \ell(y) - \ell(w) \mid w = xy, x^2 = y^2 = 1\}$. When W is finite we may also define $E(w)$, the reflection excess of w. The main result established here is that if W is finite and X is a W-conjugacy class, then there exists $w \in X$ such that w has minimal length in X and $e(w) = 0 = E(w)$. (MSC2000: 20F55)

arXiv:1405.2700v1 [math.GR] 12 May 2014

1 Introduction

Suppose that W is a Coxeter group and let $\mathcal{W}$ denote the set of strongly real elements of W. So

$$\mathcal{W} = \{w \in W \mid w = xy \text{ where } x, y \in W \text{ and } x^2 = y^2 = 1\}.$$

Let $w \in \mathcal{W}$. Then the *excess* of w, $e(w)$, is defined by

$$e(w) = \min\{\ell(x) + \ell(y) - \ell(w) \mid w = xy, x^2 = y^2 = 1\}.$$

In [7] it is shown that any element $w \in \mathcal{W}$ is W-conjugate to an element whose excess equals zero. Or, in other words, w is W-conjugate to an element xy where x and y are involutions or the identity element and $\ell(xy) = \ell(x) + \ell(y)$. The present paper explores this theme further in the case when W is finite. When W is finite it is well-known that $W = \mathcal{W}$. From Carter's seminal paper [3], it follows that, in fact, each $w \in W$ can be expressed in the form $w = xy$ where $x^2 = y^2 = 1$ and

$$V_{-1}(x) \cap V_{-1}(y) = \{0\}.$$

Here V is a reflection module for W (to be defined in Section 2) and $V_\lambda(x)$ denotes the λ-eigenspace of x on V. We note [3] only considers the Weyl groups – for a proof covering all finite Coxeter groups see Lemma 2.4. Writing $L(w)$ for the reflection length of w, this means that

$$L(w) = L(x) + L(y).$$

With this in mind we define the *reflection excess* of w, denoted $E(w)$, to be

$$E(w) = \min\{\ell(x) + \ell(y) - \ell(w) \mid w = xy, x^2 = y^2 = 1, L(w) = L(x) + L(y)\}.$$

It is clear that $E(w) \geq e(w)$. However, it is not always the case that $E(w) = e(w)$. Consider the element $w = (145)(236)$ of Sym(6), the Coxeter group of type A_5. We have $L(w) = 4$,

*The authors wish to acknowledge partial support for this work from the Birkbeck Faculty of Social Science and the Manchester Institute for Mathematical Sciences (MIMS).

x	y	$L(x)+L(y)$	$\ell(x)+\ell(y)$
(14)(23)	(15)(26)	4	$6+12=18$
(14)(36)	(15)(23)	4	$8+8=16$
(14)(26)	(15)(36)	4	$10+10=20$
(15)(23)	(45)(26)	4	$8+8=16$
(15)(36)	(45)(23)	4	$10+2=12$
(15)(26)	(45)(36)	4	$12+6=18$
(45)(23)	(14)(26)	4	$2+10=12$
(45)(36)	(14)(23)	4	$6+6=12$
(45)(26)	(14)(36)	4	$8+8=16$
(12)(46)(35)	(13)(24)(56)	6	$5+5=10$
(13)(24)(56)	(16)(25)(34)	6	$5+15=20$
(16)(25)(34)	(12)(35)(46)	6	$15+5=20$

Table 1: $(145)(236) = xy$

$\ell(w) = 10$ and Table 1 gives the possibilities for x and y with $w = xy$. As can be seen, $E(w) = 2$ but $e(w) = 0$. Surprisingly, the difference between $E(w)$ and $e(w)$ can be arbitrarily large, as is shown in Proposition 3.3.

The main theorem in this paper is

Theorem 1.1 *Suppose that W is a finite Coxeter group. If X is a conjugacy class of W, then there exists $w \in X$ of minimal length in X such that $e(w) = E(w) = 0$.*

We do not know if it is true that for an arbitrary Coxeter group and a strongly real conjugacy class X, there exists $w \in X$ with w of minimal length in X and $e(w) = 0$. However we note that this does hold for an arbitrary Coxeter group when X is a strongly real conjugacy class whose elements have finite order. This may be seen by combining a theorem of Tits (Chap V, Section 4, Ex 2d of [1]) with Lemma 2.3 (which holds in general) and Theorem 1.1.

This paper is arranged as follows. Our next section gathers together relevant background material while reviewing much of the standard notation used for Coxeter groups. Section 3, apart from proving Proposition 3.3, focusses on the proof of Theorem 1.1. Part of the proof involves checking, with the aid of MAGMA [2], all the cuspidal classes of the exceptional finite irreducible Coxeter groups. The data resulting from these calculations is documented in the appendix. At present, when studying minimal elements in conjugacy classes, this case-by-case approach is often the best we can do – see for example Chapter 3 of Geck and Pfeiffer [6].

2 Preliminary Results and Notation

From now on we assume that W is a finite Coxeter group and quickly review standard notation and facts about such groups. So W has a presentation of the form

$$W = \langle R \mid (rs)^{m_{rs}} = 1, r, s \in R \rangle$$

where $m_{rs} = m_{sr} \in \mathbb{N}$, $m_{rr} = 1$ and $m_{rs} \geq 2$ for $r, s \in R, r \neq s$. The rank of W is $|R|$. The length of an element w of W, denoted by $\ell(w)$, is defined to be

$$\ell(w) = \begin{cases} \min\{l \mid w = r_1 \cdots r_l, r_i \in R\} \text{ if } w \neq 1 \\ 0 \text{ if } w = 1. \end{cases}$$

Let V be a real euclidean vector space with basis $\Pi = \{\alpha_r \mid r \in R\}$ and norm $||\ ||$. For $r, s \in R$

$$\langle \alpha_r, \alpha_s \rangle = -||\alpha_r||\ ||\alpha_s||\ \cos\left(\frac{\pi}{m_{rs}}\right)$$

defines a symmetric bilinear form $\langle\ ,\ \rangle$ on V. Letting $r, s \in R$ we define

$$r \cdot \alpha_s = \alpha_s - \frac{2\langle \alpha_r, \alpha_s \rangle}{\langle \alpha_r, \alpha_r \rangle} \alpha_r.$$

This then extends to an action of W on V which is both faithful and respects the bilinear form $\langle\ ,\ \rangle$ (see [8]). The elements of R act as reflections upon V and V is referred to as a reflection module for W. The subset $\Phi = \{w \cdot \alpha_r \mid r \in R, w \in W\}$ of V is the root system of W, and $\Phi^+ = \{\sum_{r \in R} \lambda_r \alpha_r \in \Phi \mid \lambda_r \geq 0 \text{ for all } r\}$ and $\Phi^- = -\Phi^+$ are, respectively, the positive and negative roots of Φ. For $w \in W$, let $N(w) = \{\alpha \in \Phi^+ \mid w \cdot \alpha \in \Phi^-\}$ – it is an important fact that $\ell(w) = |N(w)|$. In a similar vein we have the following well-known result.

Lemma 2.1 *Let $g, h \in W$. Then*

$$N(gh) = N(h) \setminus [-h^{-1}N(g)] \cup h^{-1}[N(g) \setminus N(h^{-1})].$$

Hence $\ell(gh) = \ell(g) + \ell(h) - 2 \mid N(g) \cap N(h^{-1}) \mid$.

Proof See, for example, Lemma 2.2 in [7]. □

For J a subset of R define W_J to be the subgroup generated by J. Such a subgroup of W is referred to as a standard parabolic subgroup. Standard parabolic subgroups are Coxeter groups in their own right with root system

$$\Phi_J = \{w \cdot \alpha_r \mid r \in J, w \in W_J\}$$

(see Section 5.5 of [8] for more on this). A conjugate of a standard parabolic subgroup is called a parabolic subgroup of W. Finally, a *cuspidal* element of W is an element which is not contained in any proper parabolic subgroup of W. Equivalently, an element is cuspidal if its W-conjugacy class has empty intersection with all the proper standard parabolic subgroups of W.

Theorem 2.2 *Let $0 \neq v \in V$. Then the stabilizer of v in W is a proper parabolic subgroup of W.*

Proof Consult Ch V §3 Proposition 2 of [1]. □

Lemma 2.3 *Suppose that $J \subseteq R$ and $w \in W_J$. Then*

$$\min\{\ell(h^{-1}wh) \mid h \in W_J\} = \min\{\ell(g^{-1}wg) \mid g \in W\}.$$

Proof See Lemma 3.1.14 of [6]. □

Lemma 2.4 *Suppose $w \in W$. Then there exist $x, y \in W$ with $w = xy$, $x^2 = y^2 = 1$ and $V_{-1}(x) \cap V_{-1}(y) = \{0\}$.*

Proof The proof is by induction on the rank of W, the rank 1 case being trivial. Let $x, y \in W$ be such that $w = xy$ with $x^2 = y^2 = 1$. (For W a Weyl group, this is possible by [3]. The case when W is a dihedral group is straightforward to verify while types H_3 or H_4 may be checked using MAGMA [2].) If $V_{-1}(x) \cap V_{-1}(y) = \{0\}$, we are done. So suppose $0 \neq v \in V_{-1}(x) \cap V_{-1}(y)$. By Theorem 2.2, w is contained in a proper parabolic subgroup of W. Hence w is conjugate to an element u of some proper standard parabolic subgroup W_J of W. By induction $u = ab$ for some $a, b \in W_J$ where $a^2 = b^2 = 1$ and $V_{-1}(a) \cap V_{-1}(b) = \{0\}$. The appropriate conjugates of a and b will have the same properties with respect to w, so proving the lemma. □

Lemma 2.5 *Suppose that $R = \{r_1, r_2, \ldots, r_n\}$ and set $w = r_1 r_2 \cdots r_n$. Let X denote the W-conjugacy class of w. Then*

(i) the minimal length in X is n; and

(ii) the product of $r_1, \ldots, r_n$ in any order is an element of X.

Proof See Proposition 3.1.6 of [6]. □

Note that the minimal length elements of X in Lemma 2.5 are known as the *Coxeter elements* of W. We shall use $\mathrm{Dih}(2m)$ to denote the dihedral group of order $2m$. The famous classification of irreducible finite Coxeter groups obtained by Coxeter [4] (see also [8]) states

Theorem 2.6 *An irreducible finite Coxeter group is either of type $A_n (n \geq 1)$, $B_n (n \geq 2)$, $D_n (n \geq 4)$, $\mathrm{Dih}(2m) (m \geq 5)$, E_6, E_7, E_8, F_4, H_3 or H_4.*

We next discuss concrete descriptions of the Coxeter groups of types A_n, B_n and D_n which will feature in a number of our proofs. We draw the reader's attention to the fact that when regarding W as a group of permutations or signed permutations we act on the right, as is customary for permutations. First, $W(A_n)$ may be viewed as being $\mathrm{Sym}(n+1)$ with the set of fundamental reflections given by $\{(12), (23), \ldots, (n\ n+1)\}$. The elements of $W(B_n)$ can be thought of as signed permutations of $\mathrm{Sym}(n)$. We say a cycle in an element of $W(B_n)$ is of negative sign type if it has an odd number of minus signs, and positive sign type otherwise. The set of fundamental reflections in $W(B_n)$ can be taken to be $\{(\overset{++}{12}), (\overset{++}{23}), \ldots, (n-1\ \overset{+}{n}), (\bar{n})\}$. An element w expressed as a product $g_1 g_2 \cdots g_k$ of disjoint signed cycles is *positive* if the product of all the sign types of the cycles is positive, and negative otherwise. The group $W(D_n)$ consists of all positive elements of $W(B_n)$. The fundamental reflections of $W(D_n)$ can be taken to be the set $\{(\overset{++}{12}), (\overset{++}{23}), \ldots, (n-1\ \overset{+}{n}), (n-1\ \bar{n})\}$.

Let $\{e_i\}$ be an orthonormal basis for the reflection module V. For $\sigma \in W(A_n)$, define $e_i \cdot \sigma = e_{i\sigma}$. For $1 \leq i \leq n$ we may set $\alpha_{(i\ i+1)}$ to be $e_i - e_{i+1}$. The positive roots of $W(A_n)$ are then $e_i - e_j$ for $1 \leq i < j \leq n$. The positive roots of B_n are of the form $e_i \pm e_j$ for $1 \leq i < j \leq n$ and e_i for $1 \leq i \leq n$ and, for example, $(e_1 + e_2) \cdot (\overset{-+-}{124}) = -e_2 + e_4$. The positive roots of D_n are of the form $e_i \pm e_j$ for $1 \leq i < j \leq n$. Therefore the root system of

D_n consists of the long roots of the root system for B_n. Even if w is positive, it may contain negative cycles, which we wish on occasion to consider separately, so when considering elements of $W(D_n)$ we often work in the environment of $W(B_n)$ to avoid ending up with non-group elements.

The following result is obtained from Propositions 3.4.6, 3.4.7, 3.4.11, 3.4.12 of [6]. The length of minimal length elements in conjugacy classes is not given explicitly, but expressions for elements of minimal length are given in Section 3.4.2 and from these the length can be easily calculated.

Proposition 2.7 *Let W be of type B_n or D_n. Then the following hold.*

(i) Conjugacy classes in W are parameterized by signed cycle type, with one class for each signed cycle type except in the case where all cycles are of even length and positive, and W is of type D_n. In this case there are two conjugacy classes, which can be interchanged by the length-preserving graph automorphism.

(ii) Cuspidal conjugacy classes in W are those whose signed cycle type consists only of negative cycles.

(iii) Each cuspidal conjugacy class X corresponds to a non-increasing partition

$$\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_k)$$

of n, where each λ_i is the length of a negative cycle of $w \in X$. Let $\mu_i = \sum_{j=i+1}^{k} \lambda_i$. The minimal length of an element of X is

$$\begin{cases} \sum_{i=1}^{k} (\lambda_i + 2\mu_i) & \text{if } W = W(B_n) \\ \sum_{i=1}^{k} (\lambda_i - 1 + 2\mu_i) & \text{if } W = W(D_n) \end{cases}.$$

We close this section with a few pieces of notation. For a subset X of W we define $\ell_{\min}(X) = \min\{\ell(x) \mid x \in X\}$. The longest element of W will be denoted by w_0. When $W = W(A_{n-1}) \cong \mathrm{Sym}(n)$, W acts in the usual way on the set $\Omega = \{1, 2, \ldots, n\}$. In this case, for $w \in W$, the *support* of w, $\mathrm{supp}(w)$, is given by $\mathrm{supp}(w) = \{\delta \in \Omega \mid \delta w \neq \delta\}$.

3 Minimal Length and Zero Excess and Reflection Excess

We begin with an elementary lemma.

Lemma 3.1 *For $w \in W$, the following hold.*

(i) Both $e(w)$ and $E(w)$ are non-negative and even.

(ii) If w is an involution or the identity element, then $e(w) = E(w) = 0$.

(iii) $\ell(w)$ is the sum of the lengths, $e(w)$ is the sum of the excesses, and $E(w)$ is the sum of the reflection excesses of the projections of w into the irreducible direct factors of W.

Proof For (i), suppose $x^2 = y^2 = 1$ and $xy = w$. Then, using Lemma 2.1, $\ell(w) = \ell(x) + \ell(y) - 2|N(x) \cap N(y)|$ and hence $\ell(x) + \ell(y) - \ell(w)$ is even and (i) follows. The remaining parts of the lemma are straightforward. □

In the next lemma we encounter the following two subsets of W:-

$$\mathcal{I}_w = \{x \in W \mid x^2 = 1, w^x = w^{-1}\} \text{ and}$$

$$\mathcal{J}_w = \{x \in W \mid x^2 = 1, w^x = w^{-1}, V_1(w) \subseteq V_1(x)\}.$$

Lemma 3.2 *Suppose that $w \in W$.*

(i) $\mathcal{J}_w$ is the set of x such that $w = xy$ where $x^2 = y^2 = 1$ and $L(w) = L(x) + L(y)$.

(ii) If w is cuspidal, then $\mathcal{I}_w = \mathcal{J}_w$ and, in particular, $e(w) = E(w)$.

Proof Suppose $x \in \mathcal{J}_w$. Set $y = xw$. Clearly $y^2 = 1$. Let $v \in V_{-1}(x) \cap V_{-1}(y)$. Then $y \cdot v = x \cdot v$ and hence $w \cdot v = v$. That is, $v \in V_1(w)$. But $V_1(w) \subseteq V_1(x)$ as $x \in \mathcal{J}_w$. Thus $v \in V_1(x) \cap V_{-1}(x)$ and so $v = 0$. Therefore $V_{-1}(x) \cap V_{-1}(y) = \{0\}$, whence $L(w) = L(x) + L(y)$.
On the other hand, suppose $x^2 = y^2 = 1$, $w = xy$ and $L(w) = L(x) + L(y)$. Then $V_{-1}(x) \cap V_{-1}(y) = \{0\}$. Clearly $w^x = w^{-1}$. Suppose $v \in V_1(w)$. Then $v = w \cdot v = xy \cdot v$. Hence $x \cdot v = y \cdot v$. Therefore $x \cdot v - v = y \cdot v - v$. But $x \cdot v - v \in V_{-1}(x)$ and $y \cdot v - v \in V_{-1}(y)$, which forces $x \cdot v - v = y \cdot v - v = 0$. That is, $v \in V_1(x)$. Therefore $V_1(w) \subseteq V_1(x)$ and $x \in \mathcal{J}_w$, which establishes (i).

For (ii), w being cuspidal implies, by Theorem 2.2, that $V_1(w) = 0$. So $\mathcal{I}_w = \mathcal{J}_w$, and clearly $e(w) = E(w)$.

□

We now give examples of elements in the symmetric group which have arbitrarily large reflection excess, while having zero excess, showing that these quantities can differ by arbitrarily large amounts.

Proposition 3.3 *Let $W = W(A_{n-1}) \cong \mathrm{Sym}(n)$ and assume $0 < 4k \leq n$. Define*

$$\begin{aligned} w_1 &= (1\ 4\ 6\ 8 \cdots 4k-4\ 4k-2\ \ 4k-1), \\ w_2 &= (2\ \ 4k\ \ 4k-3\ 4k-5 \cdots 7\ 5\ 3) \end{aligned}$$

and $w = w_1 w_2$. Then $e(w) = 0$ but $E(w) \geq 4(k-1)^2$.

Proof First we observe that $\mathrm{supp}(w_1) \cap \mathrm{supp}(w_2) = \emptyset$. We define two involutions x, y as follows.

$$\begin{aligned} x &= (13)(2\ 4k-1)\,(45)(67)\cdots(4k-6\ 4k-5)(4k-4\ 4k-3)\,(4k-2\ 4k) \\ y &= (12)(34)\cdots(4k-1\ 4k) \end{aligned}$$

A simple check shows $w = xy$. Now $N(y) = \{e_{2i-1} - e_{2i} | 1 \leq i \leq 2k\}$. To find $N(x) \cap N(y)$, note that

$$\begin{aligned} (e_1 - e_2) \cdot x &= e_3 - e_{4k-1} \in \Phi^+ \\ (e_3 - e_4) \cdot x &= e_1 - e_5 \in \Phi^+ \\ (e_{4k-3} - e_{4k-2}) \cdot x &= e_{4k-4} - e_{4k} \in \Phi^+ \\ (e_{4k-1} - e_{4k}) \cdot x &= e_2 - e_{4k-2} \in \Phi^+ \\ (e_{2i-1} - e_{2i}) \cdot x &= e_{2i-2} - e_{2i+1} \in \Phi^+ \ (3 \leq i \leq 2k-2). \end{aligned}$$

Hence $N(x) \cap N(y) = \emptyset$, which means $e(w) = 0$ by Lemma 2.1.
Next we consider $E(w)$. Since $w_1 = (1\ 4\ 6\ 8 \cdots 4k-2\quad 4k-1)$ and $1 < 4 < \cdots < 4k-1$, we can compare w_1 seen as an element of $\mathrm{Sym}(\mathrm{supp}(w_1))$ with the element $(1\ 2 \cdots 2k)$ as an element of $\mathrm{Sym}(2k)$. This allows us to use [Proposition 2.7(iv), [7]] and Lemma 3.2(ii) to deduce that $E(w_1) \geq E((12\cdots 2k)) = 2(k-1)^2$. This is because whenever $w_1 = \sigma_1\tau_1$ where σ_1, τ_1 are involutions in $\mathrm{Sym}(\mathrm{supp}(w_1))$, we have that

$$(N(\sigma_1) \cap N(\tau_1))\,|_{\mathrm{Sym}(\mathrm{supp}(w_1))} \subseteq N(\sigma_1) \cap N(\tau_1).$$

Similarly, by comparing $w_2 = (2\ \ 4k\quad 4k-3 \cdots 7\ 5\ 3)$ to $(1\ \ 2k\ \ 2k-1 \cdots 3\ 2) = (1\ 2 \cdots 2k)^{-1}$ we deduce that $E(w_2) \geq 2(k-1)^2$.

Suppose $w = \sigma\tau$, where σ and τ are involutions with $L(\sigma) + L(\tau) = L(w)$. Then $w^{-1} = w^\sigma = w_1^\sigma w_2^\sigma$. So, as $w^{-1} = w_1^{-1}w_2^{-1}$, we have

$$w_1^\sigma w_2^\sigma = w_1^{-1}w_2^{-1}.$$

Since $\mathrm{supp}(w_1) \cap \mathrm{supp}(w_2) = \emptyset$, it follows that either $w_1^\sigma = w_1^{-1}$ or $w_1^\sigma = w_2^{-1}$. Suppose for the moment that $w_1^\sigma = w_2^{-1}$. Then $w_1^\tau = w_2^{-1}$. In particular, both σ and τ interchange $\mathrm{supp}(w_1)$ and $\mathrm{supp}(w_2)$. Put

$$v = \sum_{i \in \mathrm{supp}(w_1)} e_i - \sum_{j \in \mathrm{supp}(w_2)} e_j.$$

Then $0 \neq v \in V_{-1}(\sigma) \cap V_{-1}(\tau)$, which contradicts the fact that $L(w) = L(\sigma) + L(\tau)$. Therefore we have $w_1^\sigma = w_1^{-1}$, and so $w_2^\sigma = w_2^{-1}$. Similarly $w_1^\tau = w_1^{-1}$ and $w_2^\tau = w_2^{-1}$. As a consequence every 2-cycle of σ and every 2-cycle of τ has its support either in $\mathrm{supp}(w_1)$ or $\mathrm{supp}(w_2)$. Therefore there are involutions $\sigma_1, \sigma_2, \tau_1$ and τ_2 such that $\sigma = \sigma_1\sigma_2$, $\tau = \tau_1\tau_2$ where $w_1 = \sigma_1\tau_1$, $w_2 = \sigma_2\tau_2$, $\mathrm{supp}(\sigma_1) \cup \mathrm{supp}(\tau_1) \subseteq \mathrm{supp}(w_1)$ and $\mathrm{supp}(\sigma_2) \cup \mathrm{supp}(\tau_2) \subseteq \mathrm{supp}(w_2)$. Any $\{i, j\} \subseteq \mathrm{supp}(w_1)$ (and hence iw_1, jw_1) will thus be fixed by σ_2 and τ_2. So if $e_i - e_j \in N(\sigma_1) \cap N(\tau_1)$, then $e_i - e_j \in N(\sigma) \cap N(\tau)$. Hence

$$(N(\sigma_1) \cap N(\tau_1)) \subseteq N(\sigma) \cap N(\tau)|_{\mathrm{Sym}(\mathrm{supp}(w_1))}.$$

Similarly

$$(N(\sigma_2) \cap N(\tau_2)) \subseteq N(\sigma) \cap N(\tau)|_{\mathrm{Sym}(\mathrm{supp}(w_2))}.$$

This implies that $E(w) \geq E(w_1) + E(w_2) \geq 4(k-1)^2$, which proves the proposition. □

Lemma 3.4 *Let W be a finite Coxeter group, and X be the conjugacy class of Coxeter elements of W. Then there exists $w \in X$ of minimal length in X such that $e(w) = E(w) = 0$.*

Proof Since the Coxeter graph of every finite Coxeter group is a forest (that is, a disjoint union of trees), the Coxeter graph of W is 2-colourable. Therefore we may divide the fundamental reflections into two sets $R = R_1 \dot\cup R_2$, such that for $r, s \in R_i$, $rs = sr$. Now let $x = \prod_{r \in R_1} r$ and $y = \prod_{r \in R_2} r$. Then $w = xy$ is an element of X of minimal length, and $e(w) = E(w) = 0$. □

We remark that if W is irreducible of rank at least two, its Coxeter graph can only be 2-coloured in one way and consequently there are exactly two Coxeter elements for which $e(w) = E(w) = 0$.

Our next proposition looks at cuspidal classes. The cuspidal classes in $W(B_n)$ are Coxeter elements of a reflection subgroup that is a direct product of Coxeter groups of type B. Therefore a cuspidal element is conjugate to a product of Coxeter elements of these factors. The construction of the elements τ_i and σ_i in the proof of Proposition 3.5 may be viewed as parallel to that of x and y in the proof of Lemma 3.4. (We remark that caution must be exercised, as positive roots in the root system of a reflection subgroup are not necessarily positive roots in the root system Φ of W.)

Proposition 3.5 *Let W be irreducible of type A_n, B_n or D_n and X be a cuspidal conjugacy class of W. Then there exists $w \in X$ such that $e(w) = E(w) = 0$ and w has minimal length in X.*

Proof Suppose $W = W(A_n)$. The only cuspidal class is the class of Coxeter elements, and so by Lemma 3.4 the proposition holds.

It remains to deal with the case where W is type B_n or type D_n and X is cuspidal. Here, by Proposition 2.7, the elements of X contain only negative cycles of lengths $\lambda_1 \geq \ldots \geq \lambda_k$, where $\sum \lambda_i = n$. We will construct an element $w \in X$ of minimal length. Let $\nu_1 = 0$ and for $2 \leq i \leq k$ let $\nu_i = \sum_{j=1}^{i-1} \lambda_i$. As in Proposition 2.7, for $1 \leq i \leq k$, let $\mu_i = \sum_{j=i+1}^{k} \lambda_i$. Note that $n = \mu_i + \lambda_i + \nu_i$. By Proposition 2.7, the minimal length of elements of X, $l_{\min}(X)$, is given by

$$l_{\min}(X) = \begin{cases} \sum_{i=1}^{k} (\lambda_i + 2\mu_i) & \text{if } W = W(B_n); \\ \sum_{i=1}^{k} (\lambda_i - 1 + 2\mu_i) & \text{if } W = W(D_n). \end{cases}$$

Now for $1 \leq i \leq k$ define two involutions, τ_i and σ_i as follows.

$$\tau_i = \begin{cases} (\nu_i \overset{+}{+} 1 \;\; \nu_i \overset{+}{+} 2)(\nu_i \overset{+}{+} 3 \;\; \nu_i \overset{+}{+} 4) \cdots (\nu_i + \overset{+}{\lambda_i} - 2 \;\; \nu_i + \overset{+}{\lambda_i} - 1)(\nu_i \overset{-}{+} \lambda_i) & \text{if } \lambda_i \text{ is odd} \\ (\nu_i \overset{+}{+} 2 \;\; \nu_i \overset{+}{+} 3)(\nu_i \overset{+}{+} 4 \;\; \nu_i \overset{+}{+} 5) \cdots (\nu_i + \overset{+}{\lambda_i} - 2 \;\; \nu_i + \overset{+}{\lambda_i} - 1)(\nu_i \overset{-}{+} \lambda_i) & \text{if } \lambda_i \text{ is even} \end{cases}$$

$$\sigma_i = \begin{cases} (\nu_i \overset{+}{+} 2 \;\; \nu_i \overset{+}{+} 3)(\nu_i \overset{+}{+} 4 \;\; \nu_i \overset{+}{+} 5) \cdots (\nu_i + \overset{+}{\lambda_i} - 1 \;\; \nu_i \overset{+}{+} \lambda_i) & \text{if } \lambda_i \text{ is odd} \\ (\nu_i \overset{+}{+} 1 \;\; \nu_i \overset{+}{+} 2)(\nu_i \overset{+}{+} 3 \;\; \nu_i \overset{+}{+} 4) \cdots (\nu_i + \overset{+}{\lambda_i} - 1 \;\; \nu_i \overset{+}{+} \lambda_i) & \text{if } \lambda_i \text{ is even} \end{cases}$$

Define $w_i = \tau_i \sigma_i$, $\tau = \tau_1 \cdots \tau_k$, $\sigma = \sigma_1 \cdots \sigma_k$ and $w = \tau\sigma = w_1 \cdots w_k$. Note that if $w \in W(D_n)$, then k is even and hence τ and σ are both elements of $W(D_n)$. If λ_i is odd, then

$$w_i = (\nu_i \overset{+}{+} 1 \;\; \nu_i \overset{+}{+} 2 \;\; \nu_i \overset{+}{+} 4 \;\; \cdots \nu_i + \overset{+}{\lambda_i} - 3 \;\; \nu_i + \overset{-}{\lambda_i} - 1 \;\; \nu_i \overset{+}{+} \lambda_i \;\; \nu_i + \overset{+}{\lambda_i} - 2 \cdots \nu_i \overset{+}{+} 5 \;\; \nu_i \overset{+}{+} 3)$$

whereas if λ_i is even, then

$$w_i = (\nu_i \overset{+}{+} 1 \;\; \nu_i \overset{+}{+} 3 \;\; \nu_i \overset{+}{+} 5 \cdots \nu_i + \overset{+}{\lambda_i} - 3 \;\; \nu_i + \overset{-}{\lambda_i} - 1 \;\; \nu_i \overset{+}{+} \lambda_i \;\; \nu_i + \overset{+}{\lambda_i} - 2 \cdots \nu_i \overset{+}{+} 4 \;\; \nu_i \overset{+}{+} 2).$$

In either case, w_i is a negative λ_i-cycle. Hence $w \in X$. We now consider $\ell(\tau)$ and $\ell(\sigma)$. Firstly, note that σ is a product of distinct mutually commuting fundamental reflections. Hence $\ell(\sigma) = \sum_{i=1}^{k} \lceil \frac{\lambda_i - 1}{2} \rceil$. Now τ can be split into a product $\tau = \tau'\tau''$ where τ' consists of the 2-cycles of τ and $\tau'' = (\bar{\nu_1})(\bar{\nu_2}) \cdots (\bar{\nu_k})$. Now $\ell(\tau') = \sum_{i=1}^{k} \lfloor \frac{\lambda_i - 1}{2} \rfloor$ and $N(\tau)$ consists of roots of the form $e_a - e_b$ where for some i, $\{a, b\} \subseteq \{\nu_i + 1, \ldots, \nu_i + \lambda_i\}$. Also it is not hard to see that, thinking of τ'' as an element of $W(B_n)$,

$$N(\tau'') = \cup_{i=1}^{k} \left[\{e_{\nu_i + \lambda_i}\} \cup \{e_{\nu_i + \lambda_i} \pm e_b : \nu_i + \lambda_i < b \leq n\} \right].$$

Therefore $N(\tau') \cap N(\tau'') = \emptyset$. Hence $\ell(\tau) = \ell(\tau') + \ell(\tau'')$. If $W = W(B_n)$ we get $\ell(\tau'') = k + 2\sum_{i=1}^{k} \mu_i$ and hence $\ell(\tau) = \sum_{i=1}^{k} \lfloor \frac{\lambda_i - 1}{2} \rfloor + k + 2\sum_{i=1}^{k} \mu_i$. If $W = W(D_n)$ then $\ell(\tau'') = 2\sum_{i=1}^{k} \mu_i$ and hence $\ell(\tau) = \sum_{i=1}^{k} \lfloor \frac{\lambda_i - 1}{2} \rfloor + 2\sum_{i=1}^{k} \mu_i$.

Now $w = \tau\sigma$, so $\ell(w) \leq \ell(\tau) + \ell(\sigma)$. If $W = W(B_n)$, then

$$\begin{aligned} \ell(w) &\leq \ell(\tau) + \ell(\sigma) \\ &= \sum_{i=1}^{k} \lfloor \frac{\lambda_i - 1}{2} \rfloor + k + 2\sum_{i=1}^{k} \mu_i + \sum_{i=1}^{k} \lceil \frac{\lambda_i - 1}{2} \rceil \\ &= \sum_{i=1}^{k} \lambda_i + 2\mu_i \\ &= l_{\min}(X) \leq \ell(w). \end{aligned}$$

A similar calculation when $W = W(D_n)$ shows again that $\ell(w) \leq \ell(\tau) + \ell(\sigma) = l_{\min}(X) \leq \ell(w)$. Therefore in each case, $\ell(w) = \ell(\tau) + \ell(\sigma) = l_{\min}(X)$. Hence $e(w) = 0$ and w has minimal length in X. Since w is cuspidal, $E(w) = e(w) = 0$ and we have therefore proved Proposition 3.5. □

Proof of Theorem 1.1 We argue by induction on the rank of W – when W has rank one the theorem clearly holds. By induction and Lemma 3.1(iii) we may suppose W is an irreducible Coxeter group. If $X \cap W_J \neq \emptyset$ for some $J \subsetneq R$, then induction and Lemma 2.3 yield the result. So we may also assume X is a cuspidal class of W. Then the cases when W is of type A_n, B_n and D_n are dealt with in Proposition 3.5, while the case when W is isomorphic to $\mathrm{Dih}(2m)$ is easy to handle. For the remaining exceptional irreducible Coxeter groups we had to resort to using MAGMA [2]. The sizes of $X^0_{\min}$, where $X^0_{\min}$ denotes the set of elements in X which have minimal length in X and excess zero, are itemized in the appendix as well as giving a representative element of this set. In all cases $X^0_{\min}$ is non-empty, whence Theorem 1.1 is proven.

□

Appendix

Tables 2–5 give cuspidal classes in exceptional Weyl groups. Each class is labeled using the system in Carter [3]. Detailed information about these classes is given in [6], Tables B.1 – B.6. For a conjugacy class X let $X_{\min}$ be the set of elements of minimal length in X and $X^0_{\min}$ be the set of $w \in X_{\min}$ for which $e(x) = E(x) = 0$. For each class X we give $|X_{\min}|$, $|X^0_{\min}|$ and a representative $w \in X^0_{\min}$. Tables 6 and 7 give cuspidal classes in H_3 and H_4. For these tables the label of the class is its address in the CHEVIE list [5]. See also [6]. We have included Coxeter elements for completeness, even though they are covered by Lemma 3.4. And finally, to improve readability, we write the element, say, $r_1 r_3 r_2 r_4$ as 1324.

Class X	$\lvert X_{\min}\rvert$	$\lvert X^0_{\min}\rvert$	$w \in X^0_{\min}$
F_4	8	2	1324
B_4	14	2	124323
$F_4(a_1)$	16	16	12132343
D_4	8	6	1232343234
$C_3 + A_1$	8	6	1213213234
$D_4(a_1)$	12	12	121321343234
$A_3 + \tilde{A}_1$	16	12	12132132343234
$A_2 + \tilde{A}_2$	16	16	1213213432132343
$4A_1$	1	1	w_0

Table 2: Cuspidal Classes in F_4

Class X	$\lvert X_{\min}\rvert$	$\lvert X^0_{\min}\rvert$	$w \in X^0_{\min}$
E_6	32	2	142365
$E_6(a_1)$	80	4	12542346
$E_6(a_2)$	144	36	231423154654
$A_5 + A_1$	48	10	23423454231465
$3A_2$	80	80	123142314542314565423456

Table 3: Cuspidal Classes in E_6

Class X	$\lvert X_{\min}\rvert$	$\lvert X^0_{\min}\rvert$	$w \in X^0_{\min}$
E_7	64	2	1423657
$E_7(a_1)$	160	4	125423476
$E_7(a_2)$	280	18	12465423457
$E_7(a_3)$	366	40	1231542365476
$D_6 + A_1$	96	10	123423465423457
A_7	316	58	12365423476542345
$E_7(a_4)$	800	422	234354231435465765431
$D_6(a_2) + A_1$	708	190	13542345654231435765423
$A_5 + A_2$	420	194	12342314365423145765423 45
$D_4 + 3A_1$	32	20	123423454234565423456765423 4567
$2A_3 + A_1$	360	326	42345436542314354265431765423 4567
$7A_1$	1	1	w_0

Table 4: Cuspidal Classes in E_7

Class X	$\lvert X_{\min}\rvert$	$\lvert X^0_{\min}\rvert$	$w \in X^0_{\min}$
E_8	128	2	14682357
$E_8(a_1)$	320	4	1425423768
$E_8(a_2)$	624	10	435423145768
$E_8(a_4)$	732	40	12316542376548
$E_8(a_5)$	1516	66	1231654237687654
$E_7 + A_1$	192	10	2423454231437658
D_8	852	18	242345423165438765
$E_8(a_3)$	2696	238	23423546542314768765
$D_8(a_1)$	2040	178	1245423654765423876543
$E_8(a_7)$	2360	316	3542314654231765423148
$E_8(a_6)$	3370	422	23542314354676542314587 6
$E_7(a_2) + A_1$	1758	100	1342345423143542654238 76
$E_6 + A_2$	840	194	1234542314354654231435 4876
$D_8(a_2)$	4996	362	3142354765423458765423 1456
A_8	2816	592	1234231436542314765876542345
$D_8(a_3)$	7748	910	2314231454236576542314354876 54
$D_6 + 2A_1$	256	48	3435423143542654234568765423 4567
$A_7 + A_1$	2080	422	2314254236542345676542314354265478
$E_8(a_8)$	4480	4480	1231431542345654231456765423143546787654
$E_7(a_4) + A_1$	11592	5158	1234354234565423176542314354265437 65876543
$2D_4$	4070	1262	1242542345654276542345678765423143542654376 8
$E_6(a_2) + A_2$	16374	7438	3423145423465423143546576542876542314354265 7
$A_5 + A_2 + A_1$	3752	1910	3142345423654234567654234567876542314354265478
$D_5(a_1) + A_3$	15134	4900	4254231435465423143542678765423143542654317687
$2A_4$	7952	4058	1345423143542654231435467654231435678765423456 78
$2D_4(a_1)$	15120	15120	1423542654231765423143542654876542314354265431 76· ·542348765423
$D_4 + 4A_1$	56	42	1242315423143654231435427654231435426543187654 23· ·1435426543176543
$2A_3 + 2A_1$	1260	1090	1231423154316547654231435426543876542314354265 43· ·176542345876542345
$4A_2$	4480	4480	1231423145423143542654376542314354265431768765 42· ·3143542654317654234567876542345 6
$8A_1$	1	1	w_0

Table 5: Cuspidal Classes in E_8

Class X	$\lvert X_{\min}\rvert$	$\lvert X^0_{\min}\rvert$	$w \in X^0_{\min}$
6	4	2	312
8	6	4	21231
9	6	6	121232123
10	1	1	w_0

Table 6: Cuspidal Classes in H_3

Class X	$\lvert X_{\min}\rvert$	$\lvert X^0_{\min}\rvert$	$w \in X^0_{\min}$
11	8	2	1324
14	12	4	212431
15	18	8	32434121
16	22	10	2321234121
18	24	24	121213212343
19	34	26	32121321234121
21	12	10	2132121321234121
22	24	20	2321234321234121
23	38	30	213212134321234121
24	40	40	12121321213214321234
25	36	28	2132121321234321234121
26	24	24	121213212134321213212343
27	56	52	121232121321234321213212343
28	40	38	12121321234321213212343212343
29	60	60	121213212132124321213214321234
30	24	24	12121321213243212132123432121321234
31	36	34	121213212132123432121321234321213212343
32	40	40	1212132121321243212132124321213214321234
33	24	24	12121321213214321213212343212132123432121321234
34	1		w_0

Table 7: Cuspidal Classes in H_4

References

[1] N. Bourbaki. *Lie Groups and Lie Algebras Ch. 4–6* Springer-Verlag (2002).

[2] J.J. Cannon and C. Playoust. *An Introduction to Algebraic Programming with* Magma [draft], Springer-Verlag (1997).

[3] R.W. Carter. *Conjugacy Classes in the Weyl Group*, Compositio. Math. **25**, Facs. 1 (1972), 1–59.

[4] H.S.M. Coxeter. *The complete enumeration of finite groups of the form* $R_i^2 = (R_i R_j)^{k_{ij}} = 1$, J. London Math. Soc. 10 (1935), 21–25.

[5] M. Geck, G. Hiss, F. Lübeck, G.Malle, G.Pfeiffer. *CHEVIE – A system for computing and processing generic character tables*, Appl. Algebra Engrg. Comm. Comput. 7 (1996), 175–210.

[6] M. Geck and G. Pfeiffer. *Characters of Finite Coxeter Groups and Iwahori-Hecke Algebras*, LMS Monographs, New Series 21, (2000).

[7] S.B. Hart and P.J. Rowley. *Involution Products in Coxeter Groups*, J. Group Theory 14 (2011), no.2, 251 – 259.

[8] J.E. Humphreys. *Reflection Groups and Coxeter Groups*, Cambridge Studies in Advanced Mathematics, 29 (1990).